

The galactic disk thickness and the mass of the spherical component

D.Bizyaev^{1,2}, A.Khoperskov³, and N.Tiurina²

ABSTRACT

We present results of numerical modeling made for the galactic stellar disk embedded in the spherical halo. The non-linear dynamics of bending instabilities developed in the disk is studied. The axisymmetrical bending mode is considered as a main factor increasing the disk thickness. The model dependence between the disk thickness and the halo to disk mass ratio M_h/M_d is inferred. The modeling results enable us to estimate the dark halo mass for several edge-on galaxies. The obtained values of M_h/M_d for the galaxies considered are of order 1 - 2.

MODELING

The dynamic modeling is based on numerical integration of the motion equations for N gravitationally interacting particles [1]. The system of collisionless particles forms the disk and bulge (if any) in its central part and is embedded into the dark halo. The large-scale bending instabilities in stellar disk are considered as one of the main factor regulating the disk thickness ([1],[2],[3]).

Model parameters:

- Spherical to disk mass ratio M_s/M_d
- radial scale of the halo,
- the disk central surface density Σ .

Initial conditions

The initial disk is:

- axisymmetrical,

¹University of Texas at El Paso, US

²Sternberg Astronomical Institute, Russia

³Dept. of Physics, Volgograd State University, Russia

- in an equilibrium along the radial and vertical directions,
- gravitationally stable in the disk plane

We COMPARE the output model parameters with the observed

- rotation curve,
- the scale length h and scale height z of stellar disk
- the radial velocity dispersion (if available)

for edge-on galaxies (see Table 1). As a result, we obtain the parameters of the disk and dark halo (see Table 2). Two interesting relations between the stellar disk thickness and defined parameters are shown in Fig 1 and 2.

Table 1: Assumed parameters of the galaxies

Name	D, Mpc	h , kpc	z , kpc
UGC 6080	32.3	2.9	0.69
NGC 4738	63.6	4.7	1.30
UGC 9556	30.6	3.6	0.51
UGC 9422	45.6	3.5	0.80
NGC 5170	20.0	6.8	0.82
UGC 8286	4.8	2.0	0.26
UGC 7321	10.0	2.1	0.17
NGC 891	9.5	4.9	0.99

Table 2: Parameters obtained with the help of N-body modeling

Name	Σ , $M_{\odot}/pc^2$	M_s/M_d
UGC 6080	700	1.0
NGC 4738	970	0.5
UGC 9556	510	1.1
UGC 9422	1000	0.8
NGC 5170	460	2.2
UGC 8286	200	1.6
UGC 7321	310	2.2
NGC 891	530	1.0

CONCLUSIONS

The parameters of stellar disks and dark halo are found for the sample of eight edge-on spiral galaxies with the help of numerical N-body modeling. The bending instabilities are considered as a main factor increasing the disk thickness.

The relation between the stellar disk thickness and the relative mass of dark halo is shown for the galaxies studied. Since all but the one galaxies are bulgeless, the spherical component mass reveals the dark halo mass in our galaxies. The relation between the disk central surface density and its relative thickness z/h is presented.

DB is supported by the NASA/JPL grant NRA 99-04-OSS-058. The project is partially supported by RFBR through the grant 01-02-17597

REFERENCES

- [1] Mikhailova E., Khoperskov A., Sharpak S., 2000, in conf. proc. "Stellar Dynamics: from classic to modern", ed. by Ossipkov & Nikiforov, p.147
- [2] Hunter C., Toomre A., 1969, ApJ, 155, 747
- [3] Polyachenko V., Shukhman I., 1979, AZh, 56, 724
- [4] Vandervoort P., 1991, ApJ, 377, 49
- [5] Bizyaev D., Kajsin S., 2003, AAS 201 146.05, astro-ph/0306190




Fig. 1.— The dependence between the spherical to disk mass ratio M_s/M_d and the relative stellar disk thickness z/h is shown for our sample of galaxies by triangles. The solid line indicates the same dependence made with the help of N-body modeling (taken from [1]). Since all (except two) galaxies studied are bulgeless, M_s/M_d reveals the dark halo to disk mass ratio M_h/M_d .




Fig. 2.— The relation between the central surface density of stellar disk Σ and its relative thickness z/h is shown by the triangles for our galaxies. The squares show the same parameters taken for the galaxies from [5]. The values of Σ for those galaxies is obtained using a uniform mass to light value to coincide with the triangles.